

MEMOIRS OF THE QUEENSLAND MUSEUM

BRISBANE

© Queensland Museum
PO Box 3300, South Brisbane 4101, Australia
Phone 06 7 3840 7555
Fax 06 7 3846 1226
Email qmlib@qm.qld.gov.au
Website www.qm.qld.gov.au

National Library of Australia card number
ISSN 0079-8835

NOTE

Papers published in this volume and in all previous volumes of the *Memoirs of the Queensland Museum* maybe reproduced for scientific research, individual study or other educational purposes. Properly acknowledged quotations may be made but queries regarding the republication of any papers should be addressed to the Editor in Chief. Copies of the journal can be purchased from the Queensland Museum Shop.

A Guide to Authors is displayed at the Queensland Museum web site

A Queensland Government Project
Typeset at the Queensland Museum

A SPECIMEN RECORD OF WHITE-BELLIED STORM-PETREL FROM FRASER ISLAND, SOUTHEAST QUEENSLAND. *Memoirs of the Queensland Museum* 48(1): 123-124. 2002.- The White-bellied Storm-petrel (*Fregetta grallaria* Vieillot, 1817) lives in Southern Hemisphere subantarctic, subtropical and tropical waters (Marchant & Higgins, 1990). Four subspecies breed on isolated groups of islands and differ primarily in morphometrics (Table 1). The nominate subspecies breeds on Admiralty Island (Roach), Lord Howe Is and Kermadec Group (Macauley and Curtis Is.) and is the closest known form to E Australia. *F. g. leucogaster* is of similar size to *F. g. grallaria* and breeds on Tristan da Cunha Group, Ill St Paul and Amsterdam Is (Elliot, 1957; Jouventin, 1994.). *F. g. segethi* is small and breeds on Juan Fernandez Is (Marchant & Higgins, 1990). *F. g. titan* is the largest subspecies, breeding on central Pacific Rapa Is and the Austral Group (Murphy, 1928).

Breeding biology is poorly known; at Lord Howe Is. Egg-laying begins in late January- February (Hindwood, 1940) with offspring fledging in May (Mathews, 1928). *F. g. leucogaster* (Marchant & Higgins, 1990) and *F. g. segethi* (Bent, 1922) are also summer or late summer breeders. After breeding they disperse and become largely pelagic in subtropical and tropical waters feeding on cephalopods and crustaceans (Marchant & Higgins, 1990).

Status of the White-bellied Storm-petrel in Australia is poorly known; sight-records from the Coral Sea (Stokes & Corben, 1985), Tasman Sea (Norris, 1967), offshore SE Qld and N NSW (Marchant & Higgins, 1990) and inshore at Point Lookout, North Stradbroke Is. (Palliser, 1985); and a beach-washed specimen at Eaglehawk Neck, Tasmania in 1928 (Fletcher, 1947).

We describe a bird from 2km S of Poyungan Rocks, Fraser Island (25°25'S, 153°10'E) on 24 March 1999. The intermediate morph in good condition, Queensland Museum specimen O31230 has uniform sooty black upperparts, with

white uppertail coverts producing a sharply demarked white band above the base of the tail. Upperwings including primaries uniform sooty black; secondaries dark sooty brown. Most feathers on upperparts and wings, including the main flight feather tracts, in advanced stage of wear; some feathers on the mantle black, with white tipping (i.e. freshly moulted) indicating beginning of a body moult.

Head, chin, throat and upper-breast grey-brown shading to a white lower breast and abdomen (Figs 1, 2). Chin and throat feathers uniform grey-brown to their bases. Flanks white with short dark barring and mottling; on the right hand side a dark mark from the flanks separating white uppertail coverts from the underbody. Lateral undertail coverts black with white tipping to the feathers, enclosing the white central under tail covers having broad dark bars; this dark barring and mottling indicating an intermediate phase bird. Undertail sooty dark brown. Undersurface of the primaries and secondaries sooty dark brown, with the underwing lining white. Underwing with a broad but irregular shaped blackish leading edge enclosing the white lining.

Bill black, with a distinct downward curve towards the tip. Legs and feet black; toes not extending beyond tip of the tail.

Discussion

The White-bellied Storm-petrel specimen found on Fraser Island is clearly a bird of the nominate sub-species, with all except the tail measurements falling within this subspecies range (Table 1). This measurement of the tail was taken by Queensland Museum staff and could not be remeasured due to specimen preparation. If accurate, however this tail measurement is only just out-side the range derived from the data quoted for the subspecies by Marchant & Higgins 1990, and should not change any conclusion of specific identification of the specimen. Intermediate morph White-bellied Storm-Petrels are known only from the nominate subspecies, with the Lord Howe Island Group being the only polymorphic population. On the Lord Howe Island



FIG. 1. Dorsal view of QMO31230.



FIG. 2. Ventral view of QMO31230.

Group, the intermediate morph is uncommon, with the polymorphic ratio of light: intermediate: dark birds at 1:4:5 (van Tets & Fullagar, 1984).

Although occasionally recorded in pelagic waters on the east coast of Australia (reviewed in Marchant & Higgins, 1990), the White-bellied Storm-petrel reported here is the first substantiated specimen for mainland Australia.

Literature Cited

- BENT, A.C. 1922. Life histories of North America Petrels, Pelicans, and their Allies, Order Tubinares and Order Steganopodes. Bulletin U.S. National Museum 121:1-335.
- HINDWOOD, K.A. 1940. The birds of Lord Howe Island. *Emu* 40: 1-86.
- ELLIOTT, H.F.I. 1957. A contribution to the ornithology of the Tristan Da Cunha Group. *Ibis*, 99: 545-586.
- FLETCHER, J.A. 1947. Stray feathers: Tasmanian Notes. *Emu* 46: 394-395
- MARCHANT, S. & HIGGINS, P.J. 1990. (eds) Handbook of Australian, New Zealand & Antarctic Birds: Vol. 1 Ratites to Ducks, Part A. (Oxford University Press: Melbourne).
- MATHEWS, G.M. 1928. The birds of Norfolk and Lord Howe Islands and the Australian South Polar Quadrant with additions to the

'Birds of Australia'. Pp.139. (Witherby: London).

- MURPHY, R.C. 1928. Birds collected during the Whitney South Sea expedition IV. *American Museum Novitates* 322: 1-5.
- NORRIS, A.Y. 1967. Seabirds observations from the south-west Pacific in the southern winter. *Emu* 67: 33-54.
- PALLISER, T. 1985. The Queensland Ornithological Society Bird Report, 1984. *Sunbird* 15: 45-70.
- JOUVENTIN, P. 1994. Les Populations d'oiseaux marins des T.A.A.F.: Resume de 20 Annees de Recherche. *Alauda* 62: 44-47.
- STOKES, T. & CORBEN, C. 1985. A survey of pelagic birds in the western Coral Sea and Great Barrier Reef. *Corella* 9: 25-29.
- VAN TETS, G.F. & FULLAGAR, P.J. 1984. The status of Australian seabirds. *Australian Seabird Group Newsletter*, 20:1-25.
- Rod Hobson, Queensland Parks and Wildlife Service, Fraser Island, PMB 10, MS 2173 Rainbow Beach 4581, Australia, Present Address Queensland Parks and Wildlife Service, PO Box 731, Toowoomba 4350, Australia; David Stewart, Queensland Parks and Wildlife Service, Conservation Resource Unit, PO Box 42, Kenmore 4069, Australia; received 24 May 2001.*

TABLE 1. Morphometrics of the four subspecies of White-tailed Storm-Petrels and the specimen from Fraser Island.

	<i>grallaria</i>	<i>leucogaster</i>	<i>segethi</i>	<i>titan</i>	QMO31230
Culmen	13.0-15.0	14.2-17.0	12.6-14.0	15.0-16.3	13.5
Bill depth					5.8
Bill width					5.7
Wing	159-176	158-182	146-163	177-188	175
Tail	66-82	71-90	71-77	78-89	84
Tarsus	35.1-39.6	37-43	33-37	39-43	36
Mid toe	20.2-23.4	22-30	20-22.6	23.6-27.0	21