

VOLUME 51
PART 1

MEMOIRS
OF THE
QUEENSLAND MUSEUM

BRISBANE
31 MAY 2005

© Queensland Museum

PO Box 3300, South Brisbane 4101, Australia
Phone 06 7 3840 7555
Fax 06 7 3846 1226
Email qmlib@qm.qld.gov.au
Website www.qmuseum.qld.gov.au

National Library of Australia card number
ISSN 0079-8835

NOTE

Papers published in this volume and in all previous volumes of the *Memoirs of the Queensland Museum* may be reproduced for scientific research, individual study or other educational purposes. Properly acknowledged quotations may be made but queries regarding the republication of any papers should be addressed to the Director. Copies of the journal can be purchased from the Queensland Museum Shop.

A Guide to Authors is displayed at the Queensland Museum web site www.qmuseum.qld.gov.au/resources/resourcewelcome.html

A Queensland Government Project
Typeset at the Queensland Museum

LATE PLEISTOCENE-HOLOCENE OCCURRENCE OF *CHAEROPUS* (PERAMELIDAE) AND *MACROTIS* (THYLACOMYIDAE) FROM QUEENSLAND. *Memoirs of the Queensland Museum* 51(1): 38. Recent collections of vertebrate remains from cave systems in central-eastern and north-eastern Queensland have yielded diverse small-sized mammalian taxa. Within these faunas, four perameloid genera are present and include species of *Isoodon*, *Perameles*, *Chaeropus* and *Macrotis*. The presence of *Perameles* and *Isoodon* in these deposits is not surprising because they occur at the localities in the present day. However, Hocknull (2005) and Price (2004) report on the most easterly extent of *Perameles bougainville*, a typically arid distributed taxon. In addition, the presence of *Macrotis* and *Chaeropus* significantly increases the easterly distributions of these distinctly arid-adapted taxa. Muirhead & Godthelp (1995) reported on fossil *Chaeropus ecaudatus* from Chillagoe, northeastern Queensland, considering the age of the material to be late Pleistocene. Hocknull (2005) reported late Pleistocene *Chaeropus ecaudatus* and *Macrotis lagotis* from Mount Etna, central eastern Queensland. A new locality has yielded a specimen of *Macrotis* Thomas, 1887 and is presented herein. The locality (QML1287) is considered to be Late Pleistocene - Holocene in age based on the subfossil preservation of the excavated specimens, distinctly modern associated fauna, and the lack of associated megafauna.

Family THYLACOMYIDAE (Bensley, 1903)

Macrotis sp. (Fig 1)

Locality. QML1287, 'Dodghey's Cave', Dosey Limestone Kart, Broken River Province, 120km NW Charters Towers.

Description. QMF41971 is a left M^2 with little ware, broken root base. Max. length, 4.66mm; ant. width, 3.42mm; post. width, 2.67mm. Bulbous, sub-rectangular tooth in occlusal aspect, bearing three distinct anterior cusps (protocone, paracone and conical stylar cusp '?B'); two distinct posterior cusps (metacone and conical stylar cusp '?D'). Metaconule absent. Open, dumbbell-shaped roots. Anterior cingulum present.

Remarks. Identification of the tooth as *Macrotis* was based on the massively inflated, rectangular-ovoid occlusal crown, dumbbell-shaped molar roots, absence of the metaconule and conical stylar cusps. Muirhead (1994) listed characteristics of the dentition for both species of *Macrotis*, *M. lagotis* and *M. leucura*. Unfortunately, comparative specimens of *M. leucura* were not available for study, therefore, verification of *M. leucura* requires additional specimens and a morphometric appraisal of both species' dentition. Figure 2 illustrates the distributions (recent and fossil) of *M. lagotis*, *M. leucura* and *Chaeropus ecaudatus*. This is the second record of *Macrotis* in the fossil record of Queensland.



FIG 1. QMF41971, LM^2 in occlusal view. Scale bar = 4mm.




FIG 2. Distribution map of fossil and recent populations of A. *C. ecaudatus* (solid square), B. *M. leucura*, C. *M. lagotis* (solid circle) and *Macrotis* sp. (solid star). Recent bandicoot distributions from Strahan (1998).

The massive difference in the ranges of these three taxa when comparing late Pleistocene-Holocene to pre-European distributions indicates considerable contraction into the arid interior during the Holocene. A more detailed chronology of retraction is required to elucidate the factors influencing such a massive decline prior to European arrival, whether they be climatic, biotic and/or anthropogenic.

Literature Cited

- BENSLEY, B.A. 1903. On the evolution of the Australian Marsupialia: with remarks on the relationships of marsupials in general. The Transactions of the Royal Society of London, Zoology 9: 83-217.
- HOCKNULL, S.A. 2005. Ecological succession during the late Cainozoic of central eastern Queensland: extinction of a diverse rainforest community. *Memoirs of the Queensland Museum* 51: 39-122.
- MUIRHEAD, J. 1994. Systematics, evolution and palaeobiology of recent and fossil bandicoots (Peramelemorphia: Marsupialia). PhD thesis, University of New South Wales, Sydney. Unpubl.
- MUIRHEAD, J. & GODTHELP, H. 1995. Fossil bandicoots of Chillagoe (Northeastern Queensland) and the first known specimens of the Pig-Footed Bandicoot *Chaeropus* Ogilby, 1838 from Queensland. *Australian Mammalogy* 19:73-76.
- PRICE, G. J. 2004. Fossil bandicoots (Marsupialia: Peramelidae) and environmental change during the Pleistocene on the Darling Downs, southeastern Queensland, Australia. *Journal of Systematic Palaeontology* 2(4): 347-356.
- STRAHAN, R. (ed) 1998. The Mammals of Australia. (Australian Museum and Reed Books, Sydney).
- THOMAS, O. 1887. Description of a second species of rabbit-bandicoot (*Peragale*). *Annual Magazine of Natural History* 19 (5): 397-399.
- Scott A. Hocknull, Queensland Museum, 122 Gerler Rd. Hendra, Queensland 4011, Australia ; 1 January 2005.