

Memoirs of the Queensland Museum | **Nature**

56 (2)

© Queensland Museum 2013

PO Box 3300, South Brisbane 4101, Australia
Phone 06 7 3840 7555
Fax 06 7 3846 1226
Email qmlib@qm.qld.gov.au
Website www.qm.qld.gov.au

National Library of Australia card number
ISSN 0079-8835

NOTE

Papers published in this volume and in all previous volumes of the *Memoirs of the Queensland Museum* may be reproduced for scientific research, individual study or other educational purposes. Properly acknowledged quotations may be made but queries regarding the republication of any papers should be addressed to the Director. Copies of the journal can be purchased from the Queensland Museum Shop.

A Guide to Authors is displayed at the Queensland Museum web site www.qm.qld.gov.au

A Queensland Government Project
Typeset at the Queensland Museum

FANG HISTOLOGY OF THE LATE DEVONIAN TRISTICHOPTERID *HYNERIA LINDAE* THOMSON, 1968

Memoirs of the Queensland Museum – Nature 56(2): 311–312. 2013. Tooth and fang histology in many Palaeozoic amphibians is characterised by infolding of pleats of orthodontine (plicidentine), giving the tooth or fang a labyrinthodont structure (Schultze 1969, 1970; Warren & Davey 1992; Warren & Turner 2005). Folded orthodontine also occurs in some sarcopterygian fish, including osteolepiforms, porolepiforms, and rhizodonts (Schultze 1969, 1970; Ahlberg & Johanson 1998; Warren & Turner 2005). The tooth and fang histology of many members of the Tristichopteridae Cope, 1889, a Middle to Late Devonian osteolepiform clade, has been studied previously (e.g. Vorobyeva 1959, 1962, 1977; Schultze 1969; Lelièvre & Janvier 1986; Clément 2002). Currently, two distinct folding morphologies are recognised in tristichopterids, the simpler polyplacodont type, and the more complex eusthenodont type, the latter characterised by more extensive orthodontine infolding and a pulp cavity filled with osteodontine (Schultze 1969, 1970).

The large, derived tristichopterid *Hyneria lindae* Thomson, 1968 was first described on the basis of limited material from the Upper Devonian (Famennian) Catskill Formation of Pennsylvania, United States of America, from a site now known as 'Red Hill'. Renewed excavations at the site since 1993 have yielded additional material of this taxon, which is yet to be described (Daeschler & Shubin 2007). This note describes the histology observed in a fang of *H. lindae*. Fangs are distinguished from teeth in their large size relative to adjacent teeth (Bolt & Lombard 2001), and also in their unique pattern of development and replacement: they occur in distinct pairs, with eruption and replacement occurring in an alternating pattern (Fox *et al.* 1995; Snitting 2008; Clément *et al.* 2009).

Material and Methods. A near complete (missing only the tip) fang of *H. lindae*, measuring 35 mm long with a maximum basal diameter of 15 mm, was provided to me by Dr E. B. Daeschler for examination. The isolated fang is laterally compressed, carinated both anteriorly and posteriorly, curved posteromedially and probably came from the dermopalatine or ectopterygoid (E.B. Daeschler, pers. comm. 2011). The orthodontine is well preserved in part of the fang, but has undergone significant alteration in other parts. In the poorly preserved parts the orthodontine folds have turned dark brown or been obliterated completely, leaving a crumbly brown residue which renders the fang too fragile for thin sectioning. To permit sectioning the fang was embedded in resin (Renlam 100), and two polished sections one 8 mm from the base, the other 14 mm from the base were made. The specimen, QMF 56172, is now housed in the Geosciences Collection of the Queensland Museum.

Description. Orthodontine folds are most clearly observed near the base of the fang (Fig. 1). Orthodontine is very intensely folded into both first- and second-order folds (*sensu* Vorobyeva 1962). The total number of first-order folds in the specimen cannot be exactly determined (but is estimated to be approximately 30 at the base). The number of second-order folds per first-order fold could not be determined. Folded orthodontine extends about 3 mm in toward the centre of the fang; the remaining central region consists of osteodontine, with small vascular canals still visible despite the alteration (Fig. 1A). Dentine tubules are not visible, due to the specimen being a polished section examinable only under reflected light.

Orthodontine folding is extremely tight, with individual second-order folds pressed against each other such that there is little or no room for osteodontine or bone in between. This produces a 'branched zig zag' geometry in individual folds (Warren & Turner 2005). The overall histology of the orthodontine thus presents itself as a very regular 'hexagonal' pattern, resulting from the tightly compressed second-order folds (Fig. 1B).

Comparison to other Taxa. *Hyneria lindae* displays a peculiar and unique fang histology which combines features observed in a number of different sarcopterygian groups. The pulp cavity is filled with osteodontine, a eusthenodont characteristic observed in the tristichopterids *Eusthenodon* (Schultze 1969), *Langlieria* (Clément 2002; Clément *et al.* 2009), *Platycephalichthys* and *Jarvikinia* (Vorobyeva 1959, 1962, 1977), as well as the osteolepiform *Litoptychus* (Schultze 1969). However, the hexagonal folding pattern of orthodontine in *H. lindae* is far more regular and complex than in any known tristichopterid (cf. Vorobyeva 1959, 1962, 1977; Schultze 1969; Clément 2002); the fang histology in the genera *Mandageria* and *Cabonnichthys* cannot be described from the existing material (P.E. Ahlberg, pers. comm. 2011).




FIG. 1. Histology of *Hyneria lindae* fang, showing the distinctive 'hexagonal' orthodontine folding pattern. A, microphotograph of 8 mm parbasal section of QMF 56172; scale bar is 3 mm. B, interpretive drawing of the section in A. In B, white represents more or less unaltered orthodontine, black represents natural spaces between orthodontine folds, dark grey represents areas of significant alteration, medium grey represents dissolution cavities formed possibly through diagenesis, light grey indicates more or less unaltered osteodontine and diagonal hatching represents fractures.

It is more similar to the dendrodont pattern observed in porolepiforms (Schultze 1969, 1970; Holland 2010) and in some temnospondyls (Schultze 1969; Warren & Davey 1992), although in all of these forms it is less regular and complex than observed in *H. lindae*. Additionally, there does not appear to be any bone of attachment extending between the folds in *H. lindae*, at least as can be determined from the section near the base, which is also a dendrodont characteristic (Schultze 1969, 1970). This combination of characters prevents *H. lindae* being assigned to a distinct type of fang histology, although it is most similar to the dendrodont pattern. It is clear, however, that it is not of the polyplacodont type, which within the Tristichopteridae is considered to be restricted to more primitive members, namely *Eusthenopteron*, *Tristichopterus* (Schultze 1969, 1970) and *Notorhizodon* (Young *et al.* 1992), as well as an incomplete specimen from Morocco (Lelièvre & Janvier 1986).

Acknowledgements

Dr E.B. Daeschler is thanked for the provision of the fang. I thank Drs Daeschler, G.C. Young and C.J. Burrow, and Prof. P. E. Ahlberg, for their helpful discussions, Dr M.E. Dettmann for assistance with microphotography and Dr A.G. Cook for help in obtaining resources and sectioning the fang. Drs Daeschler, Burrow, Young, Cook and A.C. Rozefelds made many suggestions which greatly improved the manuscript.

Literature Cited

- Ahlberg, P.E. & Johanson, Z. 1998. Osteolepiformes and the ancestry of tetrapods. *Nature* **395**: 792–794.
- Bolt, J.R. & Lombard, R.E. 2001. The mandible of the primitive tetrapod *Greerapereton*, and the early evolution of the tetrapod lower jaw. *Journal of Paleontology* **75**(5): 1016–1042.
- Clément, G. 2002. Large Tristichopteridae (Sarcopterygii, Tetrapodomorpha) from the Late Famennian Evieux Formation of Belgium. *Palaeontology* **45**(3): 577–593.
- Clément, G., Snitting, D. & Ahlberg, P.E. 2009. A new tristichopterid (Sarcopterygii, Tetrapodomorpha) from the Upper Famennian Evieux Formation (Upper Devonian) of Belgium. *Palaeontology* **52**(4): 823–836.
- Cope, E.D. 1889. Synopsis of the families of Vertebrata. *American Naturalist* **23**: 849–877.
- Daeschler, E.B. & Shubin, N.H. 2007. New data on *Hyneria lindae* (Sarcopterygii; Tristichopteridae) from the Late Devonian of Pennsylvania, USA. P. 65A. Abstracts of Papers from the 67th Annual Meeting of the Society of Vertebrate Paleontology, Austin, 17–20 October, 2007.
- Fox, R.C., Campbell, K.S.W., Barwick, R.E. & Long, J.A. 1995. A new osteolepiform fish from the Lower Carboniferous Raymond Formation, Drummond Basin, Queensland. *Memoirs of the Queensland Museum* **38**(1): 97–221.
- Holland, T. 2010. Upper Devonian osteichthyan remains from the Genoa River, Victoria, Australia. *Memoirs of Museum Victoria* **67**: 35–44.
- Lelièvre, H. & Janvier, P. 1986. L'Eusthénoptéridaé (Osteichthys, Sarcopterygii) du Famennien (Dévonien supérieur) du Tafilat (Maroc): nouvelle description. *Bulletin du Muséum National d'Histoire Naturelle, Paris, Section C, Série 4* **8**(3): 351–365. [In French].
- Schultze, H.-P. 1969. Die faltenzähne der rhipidistien crossopterygier der tetrapoden und die actinopterygier gattung *Lepisosteus*; nebst einer beschreibung der zahnstruktur von *Onychodus* (struniiformer crossopterygier). *Palaeontographica Italica, New Series* **35**: 63–136.
1970. Folded teeth and the monophyletic origin of Tetrapoda. *American Museum Novitates* **2408**: 1–10.
- Snitting, D. 2008. A redescription of the anatomy of the Late Devonian *Spodichthys buettleri* Jarvik, 1985 (Sarcopterygii, Tetrapodomorpha) from East Greenland. *Journal of Vertebrate Paleontology* **28**(3): 637–655.
- Thomson, K.S. 1968. A new Devonian fish (Crossopterygii: Rhipidistia) considered in relation to the origin of the Amphibia. *Postilla* **124**: 1–13.
- Vorobyeva, E.I. 1959. A new genus of crossopterygian fish *Platycephalichthys* from the Upper Devonian of the River Lovat'. *Paleontologicheskii Zhurnal* **3**: 95–106. [In Russian].
1962. Rhizodontid crossopterygian fishes from the Devonian rocks of USSR. *Trudy Paleontologicheskogo Instituta, Academia Nauk SSSR* **44**: 1–139. [In Russian].
1977. Morphology and peculiarities of evolution of crossopterygian fish. *Trudy Paleontologicheskogo Instituta, Academia Nauk SSSR* **163**: 1–239. [In Russian].
- Warren, A.A. & Turner, S. 2005. Tooth histology patterns in early tetrapods and the presence of 'dark dentine'. *Transactions of the Royal Society of Edinburgh: Earth Sciences* **96**(2): 113–130.
- Warren, A.A. & Davey, L. 1992. Folded teeth in temnospondyls – a preliminary study. *Alcheringa* **16**(2): 107–132.
- Young, G.C., Long, J.A. & Ritchie, A. 1992. Crossopterygian fishes from the Devonian of Antarctica: systematics, relationships and biogeographic significance. *Records of the Australian Museum, Supplement* **14**: 1–77.
- Peter J. Bishop Queensland Museum, Ancient Environments Program, 122 Gerler Rd, Hendra, Qld 4011, Australia. Accepted: 12 November 2012.